

PLATE VII.

	Page.
<i>Stenopora crinita</i> , Lonsdale	49
Fig. 1. Vertical peripheral section to show the elongated tear-shaped moniliform constrictions and tabulae, x. 16.	
<i>Stenopora crinita</i> , Lonsdale?	49
Fig. 2. Axial horizontal section, with thickened and unthickened primordial walls. Acanthopores highly developed, with the concentric structure of their walls apparent, and each with a central clear spot, x. 15.	
<i>Stenopora ovata</i> , Lonsdale	55
Fig. 3. Horizontal section, with thickened and unthickened walls. In the former are visible the primordial wall, structure of the secondary thickening, and young corallites at certain angles of the calices, x. 20.	
Fig. 4. Tangential section with the whole of the walls thickened; the acanthopores highly and irregularly developed, and the centre of each occupied by a dark spot; corallites very irregularly oval, x. 20.	
Fig. 5. Longitudinal section of peripheral region to show the peculiarly crowded and exceedingly moniliform condition of the constrictions and their concentrically fibrous structure, x. 25.	
Fig. 6. Longitudinal peripheral section in which the sclerenchyma has been dissolved from the monilæ, leaving them as hollow shells, x. 15.	
FruDESCENT <i>Stenopora</i>	59
Fig. 7. Tangential section, showing enormously-thickened walls and copious development of acanthopores, x. 15.	
<i>Stenopora Leichhardti</i> , N. & E.	58
Fig. 8. Slightly oblique section of the peripheral region to show the tube-like character of the acanthopores and the generally-thickened walls, x. 20.	
<i>Stenopora tasmaniensis</i> , Lonsdale	60
Fig. 9. Vertical section partly axial, partly peripheral, showing monilæ, a tabulum, and the inner margin of the greatly-thickened periphery, x. 10.	
Figs. 1 and 3 from Singleton.	
2, 4-7 " Tasmania.	
8 " Bowen River Coal-field, Queensland.	
9 " Jerringong.	