

Constituent Parts.

Oxide of Manganese,	82.00	85
Sulphur, - - -	11.50	15
Carbonic Acid, - -	5.00	
	<hr/>	<hr/>
	98	100

Klaproth, Beit. b. iii.
s. 42.

Vauquelin, Annal.
d. Mus. vi. s. 405.

Geognostic and Geographic Situations.

It is found in Cornwall; and at Nagyag in Transylvania, along with ores of tellurium, blende, copper-pyrites, compact red manganese-ore, and brown-spar.

5. Phosphate of Manganese.

Eisenpecherz, *Werner*.

Fer phosphaté, *Broch.* t. ii. p. 533.—Manganese phosphaté, *Brong.* t. ii. p. 112.—Phosphormangan, *Karsten*, Tabel. s. 72.—Manganese phosphaté ferrifere, *Haüy*, Tabl. p. 111.—Triplit, *Haus.* Handb. b. iii. s. 1079.—Phosphate of Manganese, *Aikin*, p. 72.

External Characters.

Its colour is brownish-black, sometimes inclining to clove-brown.

It occurs massive.

Internally it is shining, glistening, or glimmering, and the lustre is resinous.

The principal fracture is concealed foliated, with a threefold rectangular cleavage, in which two of the cleavages

ages