

ance of a quiescent fluid to the motion of that body, having the same velocity in the opposite direction. The determination of these impulsions being much easier than the determination of the motions communicated by the body to the particles of the fluid, this method will be followed in most of the subsequent discussions.

The general proposition already delivered is by means sufficient for explaining the various important phenomena observed in the mutual actions of solids and fluids. In particular, it gives us no assistance in ascertaining the modifications of this resistance or impulse, which depend on the shape of the body, and the inclination of its impelled or resisted surface to the direction of the motion. Sir Isaac Newton found another hypothesis necessary; namely, that the fluid should be so extremely rare, that the distance of the particles may be incomparably greater than their diameters. This additional condition is necessary for considering their actions as so many separate collisions or impulsions on a solid body. Each particle must be supposed to have abundant room to rebound, or otherwise escape, after having made its stroke, without sensibly affecting the situations and motions of the particles which have not yet made their stroke; and the motion must be so swift as not to give time for the sensible exertion of their mutual forces of attractions and repulsions.

Keeping these conditions in mind, we may proceed to determine the impulsions made by a fluid on surfaces of every kind; and the most convenient method to pursue in this determination, is to compare them all either with the impulse which the *same surface* would receive from the fluid impinging on it perpendicularly, or with the impulse which the *same stream of fluid* would make when coming perpendicularly on a surface of such extent as to occupy the whole stream.

It will greatly abbreviate language, if we make use of a few terms in an appropriated sense.