

METRICAL ANALYSIS.

THE lyric metres of the *Electra* are the following. (1) Logaoedic, based on the choree (or 'trochee'), $- \cup$, and the cyclic dactyl $\sim \cup$, which is metrically equivalent to the choree. A logaoedic verse of 4 feet (or 'tetrapody'), composed of one cyclic dactyl and three chorees, is called Glyconic; of the 'first,' 'second,' or 'third' order, according to the place of the dactyl. Glyconics occur in the first strophe of the second Stasimon. The 'Pherecratic,' a logaoedic verse of 3 feet, occurs in the third strophe of the Parodos, per. iv., v. 3. A more detailed account of logaoedic verse will be found in *O. C.* p. lviii.

(2) Choreic, based on the choree (trochee). This occurs chiefly in verses of 4 or of 6 feet, and is often used to vary logaoedics.

(3) Dactylic, esp. in the form of the rapid tetrapody (acatalectic), as used in the Parodos (first Strophe, periods i. and iv.; second str., per. ii.; and Epode, per. iv.).

(4) Dochmiac, $\cup : - - \cup | - \wedge$. Dochmiac dimeters occur in the earlier part of the $\mu\acute{\epsilon}\lambda\omicron\varsigma \acute{\alpha}\pi\omicron \sigma\kappa\eta\nu\eta\varsigma$ (1232 ff.), and in the third Stasimon. See *O. C.* p. lix.

(5) Anapaestic dimeters, with anacrusis, are used in the Parodos (third Strophe, and Epode); and without anacrusis, in the first Kommos, second Strophe, 850 ff. Like the anapaests of *Electra's* $\theta\rho\eta\nu\omicron\varsigma$ preceding the Parodos (vv. 86—120), these belong to the class which may be described as 'free' or 'melic' anapaests, in contradistinction to the march-anapaest; see W. Christ, *Metrik*, 2nd ed., § 287, p. 247. They are especially characterised by the frequency of spondees, which give a slow and solemn movement, suited to laments (whence the name 'Klaganapäste' has sometimes been applied to them); while they also admit the converse licence of resolving long syllables (cp. commentary on 88 f.).